Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity Children Aged < 5 Years (2)

Page: 1 Run Date: 8 / 9 / 05

	Sumi	mary of T	rends ir	n Growt	h and An	emia In	dicato	rs by F	Race/Eth	nicity (3)			
Race/Ethnicity and Year		E	Birthweight	(4)			Hei	ght and \	Neight			Aner	mia
		Low		High		Short Stature (5)		Over- weight (6)		≥ 2 Yrs Overweight (6)		Low H	b/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% n <u>≥</u> 95th	Number	%
White, Not Hispanic	2004 2003 2002 2001 2000 1999 1998 1997 1996 1995	17,570 16,347 15,825 13,214 17,421 19,395 21,696 23,690 28,173 30,187	8.2 7.8 7.0 6.8 7.6 8.4 7.9 8.3 8.6 10.9	9.8 10.2 10.4 10.7 11.0 11.1 10.3 10.6 10.4	78,801 77,990 79,702 74,176 90,070 95,308 109,489 119,187 125,312 130,944	5.2 4.8 4.8 5.1 5.0 5.1 4.9 5.1 5.1 5.3	6.9 7.2 7.0 7.5 7.3 7.4 7.3 7.7 7.7	12.7 12.3 12.3 11.9 12.1 11.7 11.4 10.8 10.7 11.1	28,066 27,919 28,704 26,702 34,273 36,095 42,390 45,172 44,894 45,887	15.6 15.3 15.2 15.5 14.9 14.8 15.3 14.5 14.4	13.5 13.1 13.4 12.3 12.8 12.0 11.5 10.7 10.7	39,334 37,832 38,533 35,973 46,577 51,674 60,197 64,147 72,208 75,868	12.1 12.2 11.7 12.6 11.8 11.6 12.1 12.3 13.1 13.4
Black, Not Hispanic	2004 2003 2002 2001 2000 1999 1998 1997 1996 1995	8,651 8,401 8,217 6,704 9,584 10,304 11,241 12,662 13,364 14,449	12.3 12.1 12.2 11.6 11.7 12.0 11.7 12.6 13.4	5.9 5.4 6.3 6.3 5.9 5.9 5.9 5.9 5.9	47,514 47,529 47,190 44,539 56,062 59,230 66,764 75,850 75,471 80,413	6.5 6.4 6.6 6.8 7.0 7.0 6.8 7.0 7.0	6.4 6.6 6.4 6.6 6.4 6.5 6.6 6.9 7.0 6.9	14.7 14.3 14.3 14.2 14.4 13.8 13.3 12.6 12.3 12.4	19,570 19,609 19,215 18,073 23,531 25,036 28,511 32,966 31,794 32,625	14.2 14.4 14.2 14.3 14.1 14.6 14.1 13.2 13.1 12.8	13.3 12.9 13.0 12.2 12.5 11.5 11.1 10.2 10.2	23,837 23,882 22,962 21,801 29,602 32,659 36,849 42,148 44,698 46,072	16.8 17.7 16.5 18.3 17.8 17.7 18.1 18.5 19.7 20.8



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Children Aged < 5 Years (2)

California
Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity

Page: 2 Run Date: 8/9/05

	Sumi	mary of T	rends ir	n Growt	h and An	emia In	dicato	rs by F	Race/Eth	nicity (3)			
Race/Ethnicity and Year		В	irthweight	(4)			Hei	ght and \	Weight			Anen	nia
			Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)		≥ 2 Yrs Overweight (6)		Low Hb (7)	o/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th	Number	%
Hispanic	2004 2003 2002 2001 2000 1999 1998 1997 1996 1995	123,984 115,362 112,084 93,404 114,019 112,491 114,612 123,572 135,656 134,728	7.0 6.6 6.1 6.1 6.2 6.1 6.0 6.1 6.2 7.8	8.4 8.7 9.3 9.4 9.7 9.9 9.7 9.6 9.7	591,529 588,054 583,889 546,974 616,504 616,024 648,721 678,216 664,345 651,808	4.8 4.7 4.7 4.8 4.7 4.5 4.5 4.6	4.6 4.7 4.7 4.6 4.5 4.5 4.8 4.8	17.6 17.7 17.5 17.0 17.3 16.9 16.7 16.0 15.8 16.0	237,147 242,274 237,783 221,484 258,491 264,862 275,060 285,600 272,638 255,373	16.7 16.7 16.9 16.7 16.8 16.9 16.2 16.3 16.2	18.8 19.0 18.6 17.7 17.8 17.1 16.8 16.1 15.9 16.0	378,307 374,196 363,388 340,440 412,834 429,996 450,480 469,530 483,187 461,491	14.0 13.6 13.0 13.4 13.1 12.6 12.6 13.1 13.4 14.0
American Indian/Alaskan Native	2004 2003 2002 2001 2000 1999 1998 1997 1996 1995	407 381 362 384 484 603 532 535 623 708	7.6 8.4 6.1 4.7 5.6 6.5 3.9 7.1 5.9 9.0	8.8 13.6 11.0 8.9 10.3 10.8 15.8 11.2 10.3 11.3	2,199 2,231 2,340 2,521 3,062 3,413 3,147 3,222 3,402 3,265	5.1 6.3 5.0 6.9 4.9 5.7 4.7 5.2 4.6 5.2	6.3 5.8 5.6 5.1 6.0 5.0 5.3 5.1 5.9 5.5	17.8 18.1 19.3 18.6 17.0 16.4 16.7 17.0 16.3 16.7	876 872 959 1,007 1,282 1,387 1,370 1,403 1,403 1,301	17.2 15.7 19.5 20.2 16.4 17.9 19.6 18.5 18.0 17.0	19.7 21.8 20.8 19.7 17.5 17.5 18.2 17.7 16.4 19.6	1,304 1,324 1,341 1,489 1,963 2,208 2,101 2,071 2,381 2,485	12.6 11.9 12.5 15.2 13.9 15.4 17.1 16.5 14.4 14.0



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity
Children Aged < 5 Years (2)

Page: 3 Run Date: 8/9/05

Race/Ethnicity and Year		mary of T	Birthweight			Height and Weight							
,			Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)		≥ 2 Yrs Overweight (6)		Aner Low HI (7)	b/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% <u>></u> 95th	Number	%
Asian/Pacific Islander	2004 2003 2002 2001 2000 1999 1998 1997 1996 1995	5,410 8,736 6,637 5,920 5,373 5,362 5,854 6,937 7,956 8,605	6.8 6.7 7.4 7.7 8.2 7.4 7.3 8.4 8.0 7.9	5.1 6.5 5.4 5.5 5.5 5.5 5.7 5.5 5.5 5.4	31,355 54,114 43,633 34,663 35,708 36,906 42,543 48,585 51,261 54,160	6.8 4.9 5.6 6.1 6.3 6.7 6.8 6.8 6.7 6.9	5.8 6.1 5.9 6.3 6.4 5.9 5.3 5.9 5.9	13.6 13.4 13.6 12.8 13.6 13.3 13.6 12.9 12.7 12.2	12,431 18,282 14,676 11,996 15,210 16,320 19,043 21,493 22,135 22,232	14.6 14.3 14.5 14.5 14.5 14.5 15.1 14.3 14.5	15.3 15.2 15.0 14.7 15.2 14.2 14.6 13.9 13.8 12.5	13,468 22,761 19,224 13,620 17,204 19,330 23,685 27,725 31,449 33,289	15.2 14.3 13.5 13.9 14.4 13.9 14.2 14.6 14.6
All Other/Unknown	2004 2003 2002 2001 2000 1999 1998 1997 1996 1995	20,475 17,403 14,551 11,867 14,803 14,145 14,242 14,590 15,504 16,125	7.8 7.7 7.7 7.5 8.7 7.4 7.4 7.6 8.2 10.5	8.7 8.7 9.1 8.8 8.9 9.6 9.3 8.9 8.7	114,725 101,767 94,002 79,049 89,194 87,023 88,767 88,904 85,790 82,614	5.4 5.3 5.6 5.5 5.4 5.3 5.4 5.5	6.9 7.2 7.0 6.9 7.0 6.8 6.8 7.0 6.9 7.3	13.6 13.3 13.7 13.9 13.7 13.2 13.3 12.8 12.6 12.9	39,398 35,428 32,981 26,822 31,178 31,798 31,848 31,564 30,846 29,342	15.9 15.3 15.6 15.7 15.4 15.7 15.5 14.8 14.5	15.8 15.5 15.2 14.9 15.2 14.4 14.3 13.7 13.9	47,262 40,513 37,634 32,470 42,587 46,346 48,824 46,460 48,179 45,397	14.5 13.4 13.4 14.5 13.5 14.1 13.4 14.0 14.9



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

^{*} Percentages are not calculated if < 100 records are available for analysis after exclusions.

California

Summary of Trends in Growth and Anemia Indicators by Race/Ethnicity Children Aged < 5 Years (2)

Race/Ethnicity and Year		В	irthweight	(4)		Height and Weight							nia
		Low		High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)			Low Hb	o/Hct
		Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th - <95th	% ≥95th	Number	%
Total All Race/Ethnic Groups	2004 2003 2002 2001 2000 1999 1998 1997 1996 1995	176,497 166,630 157,676 131,493 161,684 162,300 168,177 181,986 201,276 204,802	7.4 7.1 6.7 6.6 6.9 6.9 6.8 7.0 7.2 8.9	8.3 8.6 9.1 9.1 9.4 9.6 9.4 9.2 9.3 9.2	866,123 871,685 850,756 781,922 890,600 897,904 959,431 1,013,964 1,005,581 1,003,204	5.0	5.3 5.4 5.3 5.3 5.2 5.2 5.5 5.6 5.5	16.3 16.2 16.2 15.9 16.1 15.6 15.4 14.7 14.4	337,488 344,384 334,318 306,084 363,965 375,498 398,222 418,198 403,710 386,760	16.3 16.2 16.3 16.2 16.3 16.4 15.6 15.6	17.5 17.6 17.3 16.5 16.7 15.9 15.5 14.8 14.6	503,512 500,508 483,082 445,793 550,767 582,213 622,136 652,081 682,102 664,602	14.0 13.7 13.7 13.6 13.0 13.0 14.0



Page:

Run Date:

8/9/05

⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Infants born during the reporting period included in the analysis.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

⁽⁶⁾ Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁷⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date: 9 / 20 / 05

Summary of	Trends in C	Growth ar	nd Anen	nia Ind	icators by	y Race	/Ethnicity	(3)
Race/Ethnicity and Year			Heigh	nt and W	eight			
			Short Stature (4)	Under- weight (5)			Anem Low Hb	
		Number	% <5th	% <5th	% 85th- <95th	% <u>></u> 95th	(6) Number	%
White, Not Hispanic	2004	40,186	6.0	3.1	17.0	18.7	30,027	12.3
•	2003	44,445	5.7	3.1	17.3	18.2	33,380	12.8
	2002	46,664	6.0	3.1	16.7	17.7	34,688	12.4
	2001	42,827	5.9	3.2	16.7	16.8	31,892	12.3
	2000	57,387	6.0	3.2	16.2	16.4	40,809	11.7
	1999	61,869	5.9	3.0	16.0	15.4	45,794	10.8
	1998	62,261	5.6	3.0	16.2	14.1	46,901	10.8
	1997	60,022	5.8	3.4	15.1	13.1	45,163	11.1
	1996	51,988	5.8	3.5	15.0	12.7	44,214	12.0
	1995	50,239	6.3	3.6	14.7	12.1	44,618	12.8
Black, Not Hispanic	2004	40,157	3.9	2.5	17.8	20.2	30,593	21.5
	2003	40,349	4.3	2.7	16.8	19.3	30,544	22.3
	2002	38,428	4.1	2.6	17.2	18.8	28,859	21.0
	2001	35,294	4.0	2.8	16.5	17.5	26,428	21.1
	2000	47,248	4.1	2.9	16.4	16.6	33,569	21.5
	1999	49,495	4.2	3.1	16.1	16.2	35,394	20.1
	1998	47,538	4.1	3.4	15.4	14.8	35,232	19.9
	1997	47,805	4.2	3.9	15.1	13.8	35,527	21.0
	1996	41,375	4.2	3.8	14.8	13.1	35,591	21.0
	1995	39,418	4.5	3.8	14.4	13.1	33,127	22.3



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for height-for-age.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁶⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date: 9 / 20 / 05

Summary of Tren	nds in C	Growth an	d Anen	nia Ind	icators by	y Race	/Ethnicity	(3)
Race/Ethnicity and Year			Heigh	nt and W	eight			
			Short Under- Over- Stature weight weight (4) (5) (5)			Anem Low Hb		
		Number	% <5th	% <5th	% 85th- <95th	% ≥95th	(6) Number	%
Hispanic	2004	330,393	6.2	2.3	18.9	23.8	312,758	11.5
•	2003	337,291	6.1	2.3	19.4	23.2	331,214	11.5
	2002	342,337	6.5	2.4	18.8	22.2	326,786	10.8
	2001	326,131	6.7	2.4	18.5	21.2	304,949	11.4
	2000	384,954	6.8	2.4	18.7	21.3	355,925	11.7
	1999	400,540	6.6	2.4	18.5	20.6	371,107	10.9
	1998	355,388	6.6	2.3	18.3	19.4	331,814	11.1
	1997	325,912	6.8	2.6	17.7	18.2	307,259	12.0
	1996	277,803	7.4	2.6	17.5	17.5	281,848	11.9
	1995	256,470	8.1	2.7	17.5	16.9	256,833	12.4
American Indian/Alaskan Native	2004	1,283	5.4	1.9	20.3	26.2	1,038	12.5
	2003	1,340	6.9	2.5	18.3	27.0	1,141	10.4
	2002	1,455	4.9	2.1	20.5	24.8	1,239	11.3
	2001	1,406	6.0	2.3	18.7	24.7	1,154	13.0
	2000	1,968	5.5	1.8	20.5	25.9	1,622	12.7
	1999	2,212	6.3	2.3	19.3	21.9	1,725	12.2
	1998	1,917	5.9	2.9	19.5	22.2	1,554	10.4
	1997	1,805	4.9	2.4	19.4	19.4	1,391	12.7
	1996	1,643	5.6	3.0	18.8	17.2	1,510	11.3
	1995	1,378	4.1	3.0	17.4	18.9	1,487	11.8



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for height-for-age.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁶⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date: 9 / 20 / 05

Race/Ethnicity and Year			Heigh	nt and W				
			Short Stature (4)	Under- weight (5)		Over- weight (5)		nia o/Hct
		Number	% <5th	% <5th	% 85th- <95th	% ≥95th	(6) Number	%
Asian/Pacific Islander	2004	27,600	13.8	5.0	14.1	15.2	17,252	14.1
	2003	36,958	11.4	4.8	14.0	15.2	20,517	14.4
	2002	32,276	12.3	5.0	13.8	14.2	19,110	14.0
	2001	27,026	12.6	5.1	13.7	13.2	16,326	15.1
	2000	37,206	13.7	5.3	13.0	12.9	21,827	13.3
	1999	38,569	13.2	5.4	12.6	11.6	23,711	14.1
	1998	36,892	14.5	5.7	12.1	10.6	24,786	14.7
	1997	33,631	14.6	6.3	11.1	9.9	24,067	14.6
	1996	29,441	15.7	6.5	10.5	8.9	23,704	16.2
	1995	26,394	17.7	7.4	9.9	7.8	22,423	15.9
All Other/Unknown	2004	54,821	6.8	2.9	17.9	21.5	35,157	15.4
	2003	51,821	7.1	2.9	17.4	21.4	31,641	14.5
	2002	50,983	7.5	3.0	17.5	20.0	29,557	13.7
	2001	41,809	7.3	3.0	17.0	18.9	25,884	14.7
	2000	46,057	6.9	3.3	16.8	18.8	32,313	14.4
	1999	47,420	7.4	3.4	16.6	18.0	34,876	14.4
	1998	41,459	7.5	3.4	16.2	16.2	32,731	14.3
	1997	37,874	7.7	4.1	15.5	14.9	29,816	14.5
	1996	32,376	8.0	4.0	15.4	13.9	28,075	15.3
	1995	30,003	7.8	4.3	15.4	13.9	25,707	15.9



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for height-for-age.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁶⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Page: Run Date: 9 / 20 / 05

Summary of Tre	nds in C	Growth an	d Anen	nia Ind	icators b	y Race	/Ethnicity	(3)
Race/Ethnicity and Year			Heigh					
			Short Stature (4)	Under- weight (5)			Anem Low Hb	
		Number	% <5th	% <5th	% 85th- <95th	% ≥95th	Number	%
Total All Race/Ethnic Groups	2004	494,440	6.5	2.6	18.3	22.4	426,825	12.7
	2003	512,204	6.4	2.6	18.4	21.7	448,437	12.7
	2002	512,143	6.7	2.7	18.1	20.8	440,239	12.0
	2001	474,493	6.8	2.7	17.8	19.9	406,633	12.4
	2000	574,820	6.9	2.8	17.7	19.7	486,065	12.6
	1999	600,105	6.8	2.8	17.5	18.9	512,607	11.9
	1998	545,455	6.9	2.8	17.2	17.6	473,018	12.1
	1997	507,049	7.0	3.2	16.5	16.4	443,223	12.9
	1996	434,626	7.5	3.2	16.3	15.6	414,942	13.2
	1995	403,902	8.1	3.3	16.2	15.1	384,195	13.8



⁽¹⁾ Reporting period is January 1 through December 31.

⁽²⁾ Analyses based on one record per child.

⁽³⁾ Excludes records with unknown data or errors.

⁽⁴⁾ Based on 2000 CDC growth chart percentiles for height-for-age.

⁽⁵⁾ Based on 2000 CDC growth chart percentiles for BMI-for-age. 85th - < 95th percentile category identifies children at risk of overweight.

⁽⁶⁾ Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.